

Vaccination Schedule 2021

More than
10 YEARS
of Excellence

Disease	Vaccine	Birth	6 weeks	10 weeks	14 weeks	6 months	9 months	12 months	15 months	18 months	6 years	9 years	12 years
TB	BCG	●											
Polio	bOPV	●	●										
Diphtheria, Tetanus, Polio, Pertussis Haemophilus Influenzae, Hepatitis B	Hexaximi OR Infanrix Hexa		●	●	●					●			
			8 weeks ●	12 weeks ●	16 weeks ●					●			
Pneumococcal	Prevenar 13 OR Synflorix		●	●	●		●	●	Nine and a half months				
Rotavirus	Rotarix OR RotaTeq		●	●	●								
Measles	Measbio					●		●	12 months Measbio not required if giving MMR				
Measles Mumps Rubella	Priorix OR MSD OMZYTA							●	MMR at 12 months instead of Measbio		●	Give MMR and Varilrix or Onvara separately at 6 years if not using Priorix Tetra	
Chickenpox	Varilrix OR ONVARA								●		●		
Measles Mumps Rubella + Chickenpox	Priorix Tetra	Priorix Tetra can be used at 12 months instead of Measbio, MMR and Varicella. Increased risk of febrile convulsion after the first dose of Priorix Tetra						●			●		
Meningococcal Conjugate	Menactra						●		●	For individuals 2 - 55 years administer 1 dose			
Hepatitis A	AVAXIM ⁸⁰ OR Havrix Junior							●	Hep A with MMR not with Measbio		●		
Tetanus, Diphtheria Pertussis, Polio	TETRA XIM TM OR Boostrix Tetra OR ADACEL QUADRA TM								Tetraxim at 6 years or Boostrix Tetra at 6 years Boostrix Tetra can be given at 12 years if not given at 6 years. Adacel Quadra at 12 years		●		●
Tetanus, Diphtheria	Td										●		●
Human Papilloma Virus	Cervarix OR GARDASIL	Cervarix from 9 -14 years or Gardasil from 9 -13 years as 2 doses 6 months apart State programme: Cervarix to Grade 4 girls in public schools 9 -14 years. 2 doses 6 months apart										●	
Influenza	Influvac OR VaxigripTetra	From 6 months, important for children in creche, chronic illness and respiratory problems. Start before the INFLUENZA season in MAY.											
Respiratory Syncytial Virus	SYNAGIS [®] PALIVIZUMAB	RSV prophylaxis in high-risk infants-prevention of serious LRTI caused by RSV. Start January and end in May.											
Pneumococcal	PNEUMOVAX ²³ (pneumococcal vaccine, polyvalent, MSD)	Must have at least one dose of Conjugate PCV before Pneumovax 23. Pneumovax only in children older than 2 years with Immune Compromise or high risk of Pneumococcal infection. Two doses with 2nd dose 3 - 5 years after first.											

● **STATE EPI VACCINES** These vaccines are available free from Government supplied clinics

● **RECOMMENDED OPTIONAL VACCINES** Some of the vaccines in this schedule are only available from private clinics